# NOAA COASTAL MAPPING PROGRAM PROJECT COMPLETION REPORT

#### PROJECT LA0703D

## Baptiste Collette Bayou, Louisiana

#### Introduction

NOAA Coastal Mapping Program (CMP) Project LA0703D provides highly accurate digital shoreline data for Baptiste Collette Bayou extending from the Mississippi River to Breton Sound in Louisiana. Project LA0703D is a sub-project of a larger project, LA0703 that includes large portions of the Gulf Coast from Timbalier Bay to the Mississippi River delta. The Geographic Cell (GC) can be used to complement the Nautical Charting Program (NCP) as well as geographic information systems (GIS) for a variety of coastal zone management applications.

## **Project Design**

Project LA0703D was designed per a request from the Office of Coast Survey, NOAA. Based on an analysis of project requirements and results of a source data search, it was determined that CMP procedures for multiple source projects would apply for this project. Available source data deemed adequate for successful completion of this project included sources acquired in September 2007. Due to the very large extent of complex marsh shoreline in the requested area, and the general lack of significant cultural features, various automated feature extraction techniques were used in order to expedite shoreline compilation operations.

## **Field Operations**

Routine CMP field operations did not apply for this project based on the origin of the project source data.

## Aerotriangulation

The aerotriangulation (AT) task was initiated by Remote Sensing Division (RSD) personnel in October 2009 utilizing a Digital Photogrammetric Workstation (DPW), which is a configuration of computer hardware, modular software components and other associated peripheral devices. The image files were imported into SOCET SET (SS, version 5.4.1) using the DataThruWay (version 5.4.1) software extension. The import process converted the stored compressed files to the National Imagery Transmission Format (NITF version 2.1) with headers and metadata. AT procedures were accomplished using the Multi-Sensor Triangulation (MST) module of SS. The Automatic Point Measurement algorithm within MST was used to collect tie points, and a simultaneous solve adjustment was then performed. The predicted horizontal circular error, using all measured image points, was computed to be 7 meters at the 95% confidence level. Positional data for this project is referenced to the North American Datum of 1983 (NAD 83).

## Ortho-Image

Using the image positions and orientations derived from the AT process, and the Mosaic module from within SS, an orthoimage covering the area of project LA0703D was created in December

2009. A constant elevation of zero meters (as opposed to an elevation model) was utilized to rectify the data.

## Compilation

The digital compilation phase of this project was initiated by RSD in May 2010, and entails a multi-step approach utilizing both automated and manual compilation methods, as reported below.

#### **Automated Mapping Method**

All Shoreline class features present in the database were compiled using this method. The automated feature extraction (AFE) method was performed using the orthoimage, discussed above, in combination with an object-based image analysis (OBIA) approach from within the ENVI Feature Extraction (Fx) software. Fx allows the user to interactively create and classify objects, based on image rules that meet in-situ criteria, and then to run a raster-to-vector algorithm to convert the classified image to an ArcGIS polygon shapefile. Upon completion of the AFE process, the polygon shapefile was imported into ArcGIS and 1) aggregated to merge polygons separated by less than the standard minimum distance, 2) smoothed and simplified, 3) converted to a polyline format, and 4) edited to create attribute fields compatible with the RSD interim shapefile format. The interim shapefile was then imported into SS and a Feature Data Base (FDB) was created. The FDB, consisting solely of Shoreline features, was then reviewed and edited within the SS Feature Extraction module, by the original compiler, using stereo-models derived from the AT solution.

#### Manual Mapping Method

All non-Shoreline class features present in the database (i.e., Alongshore Features, Obstructions, etc.) were compiled using this method. The manual data compilation phase of this project utilized the *traditional* RSD digital mapping approach of stereoscopic interpretation integrated with "heads-up" digitizing from within the SS Feature Extraction module. The FDB created in the automated mapping method, as described above, was subsequently populated with features derived from this manual method. This served to integrate all coastal features within a single FDB.

Feature attribution was assigned in compliance with the Coastal Cartographic Object Attribute Source Table (C-COAST), which provides the definition and attribution scheme for the cartographic features pertinent to the CMP. Selected features were further modified with additional descriptive information to refine general classification. Cartographic features were compiled to meet a horizontal accuracy of 10 meters at the 95% confidence level.

Tidal information for this project was obtained from the NOS tide station Pilots Station East, located in Southwest Pass, LA. The verified tide level for the source data was approximately 0.3 meters above Mean Lower Low Water (MLLW). The height of the Mean High Water (MHW) Datum for this tide station is 0.3 meters above the MLLW Datum.

## **Quality Control / Final Review**

Quality control tasks were conducted during all phases of project completion by a senior member of the Applications Branch of RSD. The final QC review was completed in January 2011. The

review process included analysis of the aerotriangulation and image orthorectification results, and assessment of the identification and attribution of digital feature data within the GC according to image analysis and criteria defined in C-COAST. The quality control process concluded with an inspection of topological connectivity within the GC using ArcGIS 9.3 software. All project data was evaluated for compliance to CMP requirements.

Comparisons of the largest scale NOAA nautical charts with source imagery and compiled project data resulted in creation of the Chart Evaluation File (CEF). The following nautical charts were used in the comparison process:

- Chart 11353, Baptiste Collette Bayou to Mississippi R., 1:40,000 scale, 5<sup>th</sup> ed., Jan./08
- Chart 11361, Mississippi River Delta, LA, 1:80,000 scale, 74<sup>th</sup> ed., Sep./07
- Chart 11363, Chandeleur and Breton Sounds, 1:80,000 scale, 41<sup>st</sup> ed., Jun./07
- Chart 11364, Mississippi River. Venice to New Orleans, 1:80,000, 42<sup>nd</sup> ed., Sep./07

#### **End Products and Deliverables**

The following specifies the location and identification of end products generated during the completion of this project:

#### **RSD Applications Branch Archive**

- Hardcopy of the Project Completion Report (PCR)
- Page size graphic plot of GC10818 file contents, attached to PCR

#### Remote Sensing Division Electronic Data Library


- Project database
- GC10818 in shapefile format
- Digital copy of the PCR in Adobe PDF format
- CEF in shapefile format

#### **NOAA Shoreline Data Explorer**

- GC10818 in shapefile format
- Metadata file for GC10818
- Digital copy of the PCR in Adobe PDF format

#### **End of Report**

## BAPTISTE COLLETTE BAYOU LOUISIANA







LA0703D

GC10818